



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

polar circle diagram. In translating Professor Blondel's "Moteurs Synchrone," Mr. Mailloux has rendered a valuable service to English-speaking electrical engineers.

RALPH R. LAWRENCE

Storage Batteries. By HARRY W. MORSE.
New York, The Macmillan Company. 1912.

This little book of 263 pages on storage batteries is based upon lectures given by Professor Morse at Harvard University. It deals only with the theory and the characteristics of storage batteries. No attempt is made to discuss problems connected with storage-battery engineering. The first chapters are devoted to the laws underlying the action of storage cells and to the consideration of the fundamental reactions. A short discussion of the ionic theory and the energy relations involved in the action of a storage cell is included. Later chapters are given up to the operating characteristics, efficiency and capacity, and to the general principles underlying the methods of forming modern storage battery plates. The diseases and care of storage batteries are also discussed. In the last chapter a few pages are devoted to the iron-nickel-alkali cell. "Storage Batteries" is an excellent little book for any one who wishes a simple treatment of the theory, action and care of lead-lead-peroxide storage batteries.

RALPH R. LAWRENCE

SPECIAL ARTICLES

CORRELATION BETWEEN EGG-LAYING ACTIVITY AND YELLOW PIGMENT IN THE DOMESTIC FOWL¹

In the Leghorns and the so-called American breeds, such as the Plymouth Rocks, yellow, in the form of yellow fat,² is present in varying amounts in the legs and beak. In these breeds, individual birds may undergo considerable change in the amount of the yellow pigment visible. The paling or yellowing of the

¹ Paper presented before the American Society of Naturalists, Philadelphia, December 31, 1914.

² Barrows, H. R., "Histological Basis of Shank Colors in Domestic Fowl," Bull. 232, Maine Agric. Exper. Station, 1914.

legs has been attributed by poultrymen to various environmental factors. Of recent years, some individual poultrymen, however, have claimed that paling of the legs is due to heavy laying.³ The requirements of the "Standard of Perfection," which controls judges in the show room, as well as the common practise of poultry breeders, are opposed to a belief in any connection between laying and leg color. Woods⁴ under the title, "Has Leg Color Value Indicating Layers?" in the most recent discussion of the subject, concludes:

Personally we believe that, as a practical guide in the selection of heavy layers, . . . the leg color of itself has no real value.

So far as the writers are aware, no published data are available which show in how far the leg color may be of any value in selecting the laying hen, and such suggestions as have been made in this connection have confined themselves almost entirely to a consideration of the legs alone. The results tabulated in the present paper show conclusively, it is believed, that a close connection does in fact exist between the yellow pigmentation in a hen and her previous egg-laying activity, and that, in Leghorns, the color of the ear-lobes is perhaps a better criterion of laying activity than either legs or beak and is more readily recorded.

The hens investigated were in the egg-laying contest at Storrs, Conn., and were handled essentially alike. The influence of environmental factors, therefore, can be largely neglected. The amount of yellow was measured by means of the Milton Bradley color top, which, when spinning, acts as a color mixer. The top readings were taken of the White Leghorns listed in Tables I. and II. at three different periods in October.

In Table I., the records at the three different readings have been used. A bird laying on the day of record, or on a later day within the month is considered to be laying and credited

³ Rice, J. E., Circular 11, p. 42, N. Y. State Dept. of Agriculture, 1910; Barron, Tom, *Connecticut Farmer*, September 12, 1914; Circular 499, Maine Agric. Exper. Station. This is listed as an abstract of Bull. 232.

⁴ Woods, P. T., *Amer. Poultry Jour.*, p. 35, January, 1915.